

MYPN Rabbit pAb

货号: **AYP17740**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	56kDa/114kDa/145kDa
实测分子量	145kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2
细胞定位	Cytoplasm,Nucleus,myofibril,sarcomere
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-250 of human MYPN (NP_115967.2).
序列	MQDDSI EASTSISQLLRESYLAETRHRGN NERSRAEPSSNPCHFGSPSGAAEGGGQDDL PDL SAFLSQEELDES VNLAR LAINYDPLEKADETQARKRLSPDQMKHSPNLSFEPNFCQDNPRSP TSSKESPQ EAKRPQYCSETQSKK VFLNKAAD FIEEL SSLFKSHSSKRIRPRACKNHKSKLESQNKVMQENSSSFDLSERRERSVPIPIPADTRDNEVNHAEQQEAKRREAEQAA SEAAGGDT

靶点信息

研究背景	Striated muscle in vertebrates comprises large proteins which must be organized properly to contract efficiently. Z-lines in striated muscle are a sign of this organization, representing the ends of actin thin filaments, titin, nebulin or nebulin and accessory proteins required for structure and function. This gene encodes a protein which interacts with nebulin in skeletal muscle or nebulin in cardiac muscle and alpha-actinin. In addition, this gene product can interact with a protein with the I-band indicating it has a regulatory as well as structural function. Alternative splicing results in multiple transcript variants.
------	--

基因ID	84665
基因名	MYPN
Swiss	Q86TC9
别名	MYPN;CMD1DD;CMH22;MYOP;NEM11;RCM4

产品验证



Western blot analysis of MYPN expressed in HepG2 using MYPN Rabbit pAb at 1:1000. Secondary antibody : HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn