

DNAAF1 Rabbit pAb

货号: **AYP17751**

产品信息

反应	Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	47kDa/53kDa/80kDa
实测分子量	80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat testis
细胞定位	Cell projection,Cytoplasm,cilium,cytoskeleton,spindle pole
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 516-725 of human DNA AF1 (NP_848547.4).
序列	VATEGVFTVTELDGTRTEDLETIRLETKETFCIDDLPLEDDDETGKSLEDQNMCFPKIEVISSLSDDSDPELDYTSPLVLENLPTDTLSNIFAVSKDTSKAARVPFTDIFKKEAKRDLEIRKQDTKSPRPLIQELSDDEDPSGQLLMPPTCQRDAAPLTSSGDRDSDFLAASSVPPTESAATPPETCVGVAQPSQALPTWDLTAFPAKAS

靶点信息

研究背景	The protein encoded by this gene is cilium-specific and is required for the stability of the ciliary architecture. It is involved in the regulation of microtubule-based cilia and actin-based brush border microvilli. Mutations in this gene are associated with primary ciliary dyskinesia-13. Alternative splicing results in multiple transcript variants.
基因ID	123872

基因名	DNAAF1
Swiss	Q8NEP3
别名	DNAAF1;CILD13;LRRC50;ODA7;swt

产品验证



Western blot analysis of DNAAF1 expressed in Rat testis using DNAAF1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn