

— ABLYBIO, Help Your Research



UGT2B15 Rabbit pAb

货号: **AYP17787**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:100 - 1:200
理论分子量	61kDa
实测分子量	61kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,HepG2,MCF7,BT-474,LO2,mouse liver, rat liver
细胞定位	Endoplasmic reticulum membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 60-190 of human UGT2 B15 (NP_001067.2).
------	---

靶点信息

研究背景	This gene encodes a glycosyltransferase that is involved in the metabolism and elimination of toxic compounds, both endogenous and of xenobiotic origin. This gene plays a role in the regulation of estrogens and androgens. This locus is present in a cluster of similar genes and pseudogenes on chromosome 4.
基因ID	7366
基因名	UGT2B15
Swiss	P54855
别名	UGT2B15,HLUG4,UDPGT 2B8,UDPGT2B15,UDPGTH3,UGT2B8,UGT2B15 Rabbit pAb,UDP-glucuronosyltransferase 2B8,UDPGTh-3

产品验证



Western blot analysis of UGT2B15 expressed in mouse liver, rat liver, HT-29 using UGT2B15 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)