

# CD70 Rabbit pAb

货号: **AYP17799**

## 产品信息

|       |  |
|-------|--|
| 反应    | Human,Mouse,Rat  |
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  |  |
| 应用    | IF/ICC   |
| 推荐浓度  | <b>IF/ICC:</b> 1:50 - 1:200  |
| 理论分子量 | 21kDa/23kDa  |
| 实测分子量 | 21kDa/23kDa  |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  |  |
| 细胞定位  | Membrane,Single-pass type II membrane protein  |
| 纯化    | Affinity purification  |

## 抗原信息

|      |   |
|------|---|
| 抗原信息 | A synthetic peptide corresponding to a sequence within amino acids 100-193 of human CD70 (NP_001243.1). |
| 序列   | RDGIYMVHIQVTLAICSSTTASRHHPTTLAVGICSPASRSISLLRLSFHQGCTIASQRLTPLARGDTLCTNLTGTLPSRNTDETFFGVQWVRP           |

## 靶点信息

|      |  |
|------|--|
| 研究背景 | The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin synthesis. |
| 基因ID | 970  |

|       |  |
|-------|--|
| 基因名   | CD70                                   |
| Swiss | P32970                                 |
| 别名    | CD70;CD27-L;CD27L;CD27LG;TNFSF7;TNLG8A |

产品验证

实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)