

HVEM/TNFRSF14 Rabbit pAb

货号: **AYP17908**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	21kDa/30kDa
实测分子量	30kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 39-202 of human HVEM/TNFRSF14 (NP_003811.2).
序列	LPSCKEDEYPVGSECCPKCSPGYRVKEACGELTGTVCEPCPPGTIYIAHLNGLSKLQCQMCDPAMGLRASRNCSTRNAV CGCSPGHFCIVQDGDHCAACRAYATSSPGQRVQKGGTESQDTLCQNCPPGTFSPNGTLEECQHQTCSWLVTKAGAG TSSSHWV

靶点信息

研究背景	This gene encodes a member of the TNF (tumor necrosis factor) receptor superfamily. The encoded protein functions in signal transduction pathways that activate inflammatory and inhibitory T-cell immune response. It binds herpes simplex virus (HSV) viral envelope glycoprotein D (gD), mediating its entry into cells. Alternative splicing results in multiple transcript variants.
基因ID	8764

基因名	TNFRSF14
Swiss	Q92956
别名	TNFRSF14;ATAR;CD270;HVEA;HVEM;LIGHTR;TR2

产品验证



Western blot analysis of HVEM/TNFRSF14 expressed in Raji using HVEM/TNFRSF14 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn