— ABLYBIO, Help Your Research

CCL22 Rabbit pAb

货号: AYP17911

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	IHC
推荐浓度	**IHC:** 1:50 - 1:200
理论分子量	10kDa
实测分子量	10kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Macrophage,Monocyte,Dendritic cell
细胞定位	Secreted
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-93 of human CCL22 (NP_002981.2).

靶点信息

研究背景	This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for monocytes, dendritic cells, natural killer cells and for chronically activated T lymphocytes. It also displays a mild activity for primary activated T lymphocytes and has no chemoattractant activity for neutrophils, eosinophils and resting T lymphocytes. The product of this gene binds to chemokine receptor CCR4. This chemokine may play a role in the trafficking of activated T lymphocytes to inflammatory sites and other aspects of activated T lymphocyte physiology.
基因ID	6367
基因名	CCL22
Swiss	O00626 (https://www.uniprot.org/uniprotkb/O00626/entry)
别名	CCL22,A-152E5.1,ABCD-1,DC/B-CK,MDC,SCYA22,STCP-1,CCL22 Rabbit pAb,CC chemokine STCP-1,MDC(1-69),Macrophage-derived chemokine,Small-inducible cytokine A22,Stimulated T-cell chemotactic protein 1

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)