

— ABLYBIO, Help Your Research



DLC1 Rabbit pAb

货号: **AYP17927**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	50kDa/54kDa/114kDa/122kDa/126kDa/170kDa
实测分子量	171kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse lung
细胞定位	Cell junction,Cytoplasm,Membrane,Peripheral membrane protein,focal adhesion
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 204-463 of human DLC1 (NP_079043.3).
------	--

靶点信息

研究背景	This gene encodes a GTPase-activating protein (GAP) that is a member of the rhoGAP family of proteins which play a role in the regulation of small GTP-binding proteins. GAP family proteins participate in signaling pathways that regulate cell processes involved in cytoskeletal changes. This gene functions as a tumor suppressor gene in a number of common cancers, including prostate, lung, colorectal, and breast cancers. Multiple transcript variants due to alternative promoters and alternative splicing have been found for this gene.
基因ID	10395
基因名	DLC1
Swiss	Q96QB1 (https://www.uniprot.org/uniprotkb/Q96QB1/entry)
别名	DLC1,ARHGAP7,HP,STARD12,p122-RhoGAP,DLC1 Rabbit pAb,Deleted in liver cancer 1 protein,HP protein,Rho-type GTPase-activating protein 7,START domain-containing protein 12,StAR-related lipid transfer protein 12,KIAA1723

产品验证



Western blot analysis of DLC1 expressed in Mouse lung using DLC1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)