

POLR2D Rabbit pAb

货号: **AYP17947**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:10 - 1:100 IP: 1:50 - 1:200
理论分子量	16kDa
实测分子量	17kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF-7,SW480,SW620,HepG2,SK-HEPC,Mouse spleen
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-142 of human POLR2D (NP_004796.1).
序列	MAAGGSDPRAGDVEEDASQLIFPKEFETAETLLNSEVHMLLEHRKQQNESAEDEQELSEVFMKTLNNTARFSRFRKNRETIASVRSLLLQKLLHKFELACLANLCPETAESKALIPSLEGRFEDEELQQLDDIQTKRSFQY

靶点信息

研究背景	This gene encodes the fourth largest subunit of RNA polymerase II, the polymerase responsible for synthesizing messenger RNA in eukaryotes. In yeast, this polymerase subunit is associated with the polymerase under suboptimal growth conditions and may have a stress protective role. A sequence for a ribosomal ps eudogene is contained within the 3' untranslated region of the transcript from this gene.
------	---

基因ID	5433
基因名	POLR2D
Swiss	O15514
别名	POLR2D;HSRBP4;HSRPB4;RBP4;RPB16

产品验证



Western blot analysis of POLR2D expressed in Mouse spleen, MCF-7, SW480 using POLR2D Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn