

CAR/CXADR Rabbit pAb

货号: **AYP17958**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	9kDa/22kDa/28kDa/35kDa/38kDa/40kDa
实测分子量	50KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,Mouse kidney
细胞定位	Basolateral cell membrane,Cell junction,Cell membrane,Secreted,Single-pass membrane protein,Single-pass type I membrane protein,Single-pass type I membrane protein,adherens junction,tight junction
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 20-237 of human CAR/CXADR (NP_001329.1).
序列	LSITTPEEMIEKAKGETAYLPCKFTLSPEDQGPLDIEWLISPADNQKVDQVILYSGDKIYDDYYPDLKGRVHFTSNDLKSGDASINVTNLQLSDIGTYQCKVKKAPGVANKKIHLVVLVKPSGARCVDGSEEEIGSDFKIKCEPKEGSLPLQYEWQKLSDSQKMPTSWLAEMTSSVISVKNASSEYSGTYSCTVRNRVGSQCLLRLLNVPPSNKAG

靶点信息

研究背景	The protein encoded by this gene is a type I membrane receptor for group B coxsackieviruses and subgroup C adenoviruses. Several transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene are found on chromosomes 15, 18, and 21.
基因ID	1525

基因名	CXADR
Swiss	P78310
别名	CXADR;CAR;CAR4/6;HCAR

产品验证



Western blot analysis of CAR/CXADR expressed in U-251MG, Mouse kidney using CAR/CXADR Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn