

GGCX Rabbit pAb

货号: **AYP17963**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	80kDa/87kDa
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,MCF7,HeLa,Mouse liver
细胞定位	Endoplasmic reticulum membrane,Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 479-758 of human GGC X (NP_000812.2).
序列	QRIFDPRVDIVQAAWSPFQRTSWVQPLLDLSPWRAKLQEIKSSLDNHTEVVFIADFPGLHLENFVSEDLGNTSIQLLQGE VTVELVAEQKNQTLREGEKMQLPAGEYHKVYTTSPSPSCYMYVYVNTTELALQDLAYLQELKEKVENGSETGPLPELQPL LEGEVKGKGGPEPTPLVQTFLRRQQLQEIERRRNTPFHERFFRLLRKLKLYVFRSFLMTCISLRNLILGRPSLEQLAQEVTYANL RPF EAVGELNPSNTDSSHSNPPESNPDPVHSEF

靶点信息

研究背景	This gene encodes an integral membrane protein of the rough endoplasmic reticulum that carboxylates glutamate residues of vitamin K-dependent proteins to gamma carboxyl glutamate, a modification that is required for their activity. The vitamin K-dependent protein substrates have a propeptide that binds the enzyme, with carbon dioxide, dioxide, and reduced vitamin K acting as co-substrates. Vitamin K-dependent proteins affect a number of physiologic processes including blood coagulation, prevention of vascular calcification, and inflammation. Allelic variants of this gene have been associated with pseudoxanthoma elasticum-like disorder with associated multiple coagulation factor deficiency. Alternative splicing results in multiple transcript variants.
基因ID	2677
基因名	GGCX
Swiss	P38435
别名	GGCX;VKCFD1

产品验证



Western blot analysis of GGCX expressed in Mouse liver, LO2, MCF7 using GGCX Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn