

MOK Rabbit pAb

货号: **AYP17969**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	7kDa/10kDa/27kDa/44kDa/47kDa/48kDa
实测分子量	40kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-230 of human MOK (NP_055041.1).
序列	MKNYKAIGKIGEGTFSEVMKMQSLRDGNYACKQMKQRFESIEQVNNLREIQALRRLNPHPNILMLHEVVFDRKSGSLALICELMDMNIYELIRRRYPLSEKKIMHYMYQLCKSLDHIHRNGIFHRDVKPENILIKQDVLKLGDFGSCRSVYSKQPYTEYISTRWYRAPECLLTDFYTYKMDLWSAGCVFYEIASLQPLFPGVNELDQISKIHDVIGTPAQKILTKFK

靶点信息

研究背景	This gene belongs to the MAP kinase superfamily. The gene was found to be regulated by caudal type transcription factor 2 (Cdx2) protein. The encoded protein, which is localized to epithelial cells in the intestinal crypt, may play a role in growth arrest and differentiation of cells of upper crypt and lower villus regions. Multiple alternatively spliced transcript variants encoding different isoforms have been observed for this gene.
------	--

基因ID	5891
基因名	MOK
Swiss	Q9UQ07
别名	MOK;RAGE;RAGE-1;RAGE1;STK30

产品验证



Western blot analysis of MOK expressed in Mouse liver using MOK Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn