

# DCTN1 Rabbit pAb

货号: **AYP17970**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IP:</b> 1:50 - 1:100
理论分子量	126kDa/127kDa/136kDa/138kDa/140kDa/141kDa
实测分子量	150kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,HeLa,HepG2,22RV1,Jurkat,Mouse testis,Rat testis
细胞定位	Cytoplasm,cytoskeleton
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 945-1139 of human DCTN1 (NP_001128513.1).
序列	PGLVKDSPLLLQQISAMRLHISQLQHENSILKGAQMKASLASLPLHVAKLSHEGPGSELPAAGALYRKTSQLETLNQLSTH THVVDITRTSPAAKSPSAQLMEQVAQLKSLSDTVEKLVKDEVVKETVSRPGATVPTDFATFPSSAFLRAKEEQDDTVYM GKVTFSCAAGFGQRHRLVLTQEQLHQLHSRLIS

## 靶点信息

研究背景	This gene encodes the largest subunit of dynactin, a macromolecular complex consisting of 10 subunits ranging in size from 22 to 150 kD. Dynactin binds to both microtubules and cytoplasmic dynein. Dynactin is involved in a diverse array of cellular functions, including ER-to-Golgi transport, the centripetal movement of lysosomes and endosomes, spindle formation, chromosome movement, nuclear positioning, and axonogenesis. This subunit interacts with dynein intermediate chain by its domains directly binding to dynein and binds to microtubules via a highly conserved glycine-rich cytoskeleton-associated protein (CAP-Gly) domain in its N-terminus. Alternative splicing of this gene results in multiple transcript variants encoding distinct isoforms. Mutations in this gene cause distal hereditary motor neuropathy type VIIIB (HMN7B) which is also known as distal spinal and bulbar muscular atrophy (dSBMA).
基因ID	1639
基因名	DCTN1
Swiss	Q14203
别名	DCTN1;DAP-150;DP-150;P135

## 产品验证



Western blot analysis of DCTN1 expressed in Mouse testis,Rat testis,U-251MG using DCTN1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)