

# USP33 Rabbit pAb

货号: **AYP18031**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:100
理论分子量	93kDa/103kDa/106kDa
实测分子量	107KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HepG2
细胞定位	Cytoplasm,Golgi apparatus,centrosome,cytoskeleton,microtubule organizing center,perinuclear region
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 290-400 of human USP33 (NP_963920.1).
序列	QVMEVEEDPQTITTEETMEEDKSQSDVDFQSCESCNSNDRAENENGRFCFSEDNNETTMLIQDDENNSEMskdwQKEKMCNKINKVNSEGEFDKDRDSISETVLDLNNQETV

## 靶点信息

研究背景	This gene encodes a deubiquinating enzyme important in a variety of processes, including Slit-dependent cell migration and beta-2 adrenergic receptor signaling. The protein is negatively regulated through ubiquitination by von Hippel-Lindau tumor protein (VHL). Alternative splicing results in multiple transcript variants and protein isoforms.
------	--

基因ID	23032
基因名	USP33
Swiss	Q8TEY7
别名	USP33;VDU1

## 产品验证



Western blot analysis of USP33 expressed in HeLa, HepG2 using USP33 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)