

— ABLYBIO, Help Your Research



# CHRM1 Rabbit pAb

货号: **AYP18077**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	50kDa/51kDa
实测分子量	51KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,U-87MG,C6
细胞定位	Cell junction,Cell membrane,Multi-pass membrane protein,postsynaptic cell membrane,synapse
纯化	Affinity purification


## 抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 250-350 of human CHRM1 (NP_000729.2).
------	--

## 靶点信息

研究背景	The muscarinic cholinergic receptors belong to a larger family of G protein-coupled receptors. The functional diversity of these receptors is defined by the binding of acetylcholine and includes cellular responses such as adenylate cyclase inhibition, phosphoinositide degeneration, and potassium channel mediation. Muscarinic receptors influence many effects of acetylcholine in the central and peripheral nervous system. The muscarinic cholinergic receptor 1 is involved in mediation of vagally-induced bronchoconstriction and in the acid secretion of the gastrointestinal tract. The gene encoding this receptor is localized to 11q13.
基因ID	1128
基因名	CHRM1
Swiss	P11229 ( <a href="https://www.uniprot.org/uniprotkb/P11229/entry">https://www.uniprot.org/uniprotkb/P11229/entry</a> )
别名	CHRM1, HM1, M1, M1R, CHRM1 Rabbit pAb

## 产品验证



Western blot analysis of CHRM1 expressed in SH-SY5Y, U-87MG, C6 using CHRM1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)