

# ACMSD Rabbit pAb

货号: **AYP18095**

## 产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	31kDa/38kDa
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney,Mouse liver,Rat kidney
细胞定位	cytoplasm,cytosol
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-336 of human ACMSD (NP_612199.2).
序列	MKIDIHSHILPKEWPDLLKRFYGGWVQLQHHSKGEAKLLKDGKVFVRVRENCWDPEVRIEMDQKGVTVQALSTVPV MFSYWAKPEDTLNLCQLLNNDLASTVVSYPFRFVGLGTLPMQAPELAVKEMERCVKELGFPQVIGTHVNEWDLNAQEL FPVYAAAERLKCSLFVHPWDMQMDGRMAKYWLPWLVGMPAETTIAICSMIMGGVFEKFKPKLVCFAHGGGAFPFTVGRI SHGFMRPDLCAQDNPMNPKKYLGSFYTDALVHDPLSLKLLTDVIGKDKVILGTDYFPPLGELEPGKLIESMEEFDEETKN KLGAGNALAFLGLERKQFE

## 靶点信息

研究背景	The neuronal excitotoxin quinolinate is an intermediate in the de novo synthesis pathway of NAD from tryptophan, and has been implicated in the pathogenesis of several neurodegenerative disorders. Quinolinate is derived from alpha-amino-beta-carboxy-muconate-epsilon-semialdehyde (ACMS). ACMSD (ACMS decarboxylase; EC 4.1.1.45) can divert ACMS to a benign catabolite and thus prevent the accumulation of quinolinate from ACMS.
------	--

基因ID	130013
基因名	ACMSD
Swiss	Q8TDX5
别名	ACMSD

### 产品验证



Western blot analysis of ACMSD expressed in Mouse kidney, Mouse liver, Rat kidney using ACMSD Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 12 0s.

### 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)