

— ABLYBIO, Help Your Research



SSC4D Rabbit pAb

货号: **AYP18098**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	29kDa/60kDa
实测分子量	61kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,HepG2,Mouse liver,Mouse brain,Rat liver,Rat brain
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 446-575 of human SSC4 D (NP_542782.1).
------	--

靶点信息

研究背景	The scavenger receptor cysteine-rich (SRCR) superfamily is an ancient and highly conserved group of cell surface and/or secreted proteins, some of which are involved in the development of the immune system and the regulation of both innate and adaptive immune responses. Group B SRCR domains usually contain 8 regularly spaced cysteines that give rise to a well-defined intradomain disulfide-bond pattern.
基因ID	136853
基因名	SSC4D
Swiss	Q8WTU2 (https://www.uniprot.org/uniprotkb/Q8WTU2/entry)
别名	SSC4D,S4D-SRCRB,SrcRB-S4D,SrcRB4D,SSC4D Rabbit pAb,Four scavenger receptor cysteine-rich domains-containing protein

产品验证



Western blot analysis of SSC4D expressed in Mouse liver,Rat liver,U-87MG using SSC4D Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)