

PPIL4 Rabbit pAb

货号: **AYP18127**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	57kDa
实测分子量	65kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	293T, U-87MG, LO2
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 301-492 of human PPIL4 (NP_624311.1).
序列	MDNVLIDRRRIHVDFSQSVAKVKWKGKGGKYTKSDFKEYEKEQDKPPNLVLKDKVKPKQDTKYDLILDEQAEDSKSSHS HTSKKHKKKTHHCSEEKEDDYMPIKNTNQDIYREMGFGHYEEEEESWEKQKSEKRDRTQNRSRSRERDGHYSNSH KSKYQTDLYERERSKKRDRSRSPKSKDKEKSKYR

靶点信息

研究背景	This gene is a member of the cyclophilin family of peptidylprolyl isomerases. The cyclophilins are a highly conserved family, members of which play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions.
基因ID	85313

基因名	PPIL4
Swiss	Q8WUA2
别名	PPIL4;HDCME13P

产品验证



Western blot analysis of PPIL4 expressed in 293T,U-87MG,LO2 using PPIL4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn