

CD99L2 Rabbit pAb

货号: **AYP18138**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	15kDa/20kDa/22kDa/27kDa/28kDa
实测分子量	28kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Mouse brain,Mouse kidney,Mouse liver,Mouse lung,Rat pancreas
细胞定位	Cell junction,Cell membrane,Extracellular side,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 26-185 of human CD99L2 (NP_113650.2).
序列	DFDDFNLEDAVKETSSVKQPWDHTTTTTTNRPGTTRAPAKPPGSGLDLADALDDQDDGRRKPGIGGRERWNHVTTTTKRPVTTTRAPANTLGNDFDLADALDDRNRDDGRRKPIAGGGGFSDKDLEDIVGGGEYKPKDKGKGDRYGSNDDPGSGMVAEPG

靶点信息

研究背景	This gene encodes a cell-surface protein that is similar to CD99. A similar protein in mouse functions as a n adhesion molecule during leukocyte extravasation. Alternate splicing results in multiple transcript variants.
基因ID	83692

基因名	CD99L2
Swiss	Q8TCZ2
别名	CD99L2;CD99B;MIC2L1

产品验证



Western blot analysis of CD99L2 expressed in Mouse brain,Rat pancreas,HeLa using CD99L2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn