

— ABLYBIO, Help Your Research



NGLY1 Rabbit pAb

货号: **AYP18161**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	14kDa/63kDa/70kDa/72kDa/74kDa
实测分子量	74KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U266,K-562,Mouse brain,Rat brain
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human NGLY1 (NP_060767.2).
------	-------------------------------------------------------------------------------------------------------------------

靶点信息

研究背景	This gene encodes an enzyme that catalyzes hydrolysis of an N(4)-(acetyl-beta-D-glucosaminyI) asparagine residue to N-acetyl-beta-D-glucosaminyIamine and a peptide containing an aspartate residue. The encoded enzyme may play a role in the proteasome-mediated degradation of misfolded glycoproteins. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	55768
基因名	NGLY1
Swiss	Q96IV0 (https://www.uniprot.org/uniprotkb/Q96IV0/entry)
别名	NGLY1,CDDG,CDG1V,PNG1,PNGase,NGLY1 Rabbit pAb,N-glycanase 1,Peptide:N-glycanase

产品验证



Western blot analysis of NGLY1 expressed in Mouse brain,Rat brain,U266 using NGLY1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)