

RNF220 Rabbit pAb

货号: **AYP18173**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	39kDa/62kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Mouse spleen,Mouse thymus,Mouse brain,Rat testis
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 267-566 of human RNF220 (NP_060620.2).
序列	PKSLLLSASIKREGESPTASPHSSATDDLHHSDDRYQTFLRVRANRQTRLNARIGKMKRRKQDEGQREGSCMAEDDAVDIE HENNNRFEYEWCGQKRIRATTLLEGGFRGSGFIMCSGKENPDSADLDVDGDDTLEYGKPQYTEADVIPCTGEEPGEA KEREALRGAVLNGGPPSTRITPEFSKWASDEMPSTSNGESSKQEQAMQKTCKNSDIEKITEDSAVTTFEALKARVRELERQL SRGDRYKCLICMDSYSMPLTSIQCWVHCEECWLRTLGAKKLCPCNTITAPGLRRIYL

靶点信息

研究背景	E3 ubiquitin-protein ligase that promotes the ubiquitination and proteasomal degradation of SIN3B (By similarity). Independently of its E3 ligase activity, acts as a CTNNB1 stabilizer through USP7-mediated deubiquitination of CTNNB1 promoting Wnt signaling.
基因ID	55182

基因名	RNF220
Swiss	Q5VTB9
别名	RNF220;C1orf164

产品验证



Western blot analysis of RNF220 expressed in Mouse spleen, Rat testis, HeLa using RNF220 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn