

# TRIM62 Rabbit pAb

货号: **AYP18175**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	40kDa/54kDa
实测分子量	54kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,Mouse brain
细胞定位	Cytoplasm
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 120-280 of human TRIM 62 (NP_060677.2).
序列	HEQHQVTGIDDAFDELQRELKDQLQALQDSEREHTEALQLLKRQLAETKSSTKSLRRTTIGEAFERLHRLLRERQKAMLEELEADTARTLTIDIEQKVQRYSQQLRQVQEGAQILQERLAETDRHTFLAGVASLSERLKGKIHETNLTYEDFPTSKYTGPLQY

## 靶点信息

研究背景	E3 ubiquitin ligase that plays a role in antifungal immunity by mediating 'Lys-27'-linked ubiquitination of CARD9 downstream of C-type lectin receptors; leading to CARD9 activation, followed by activation of NF-kappa-B and MAP kinase p38 pathways. E3 ubiquitin ligase activity is dependent on E2 ubiquitin-conjugating enzyme UBE2D2.
基因ID	55223

基因名	TRIM62
Swiss	Q9BVG3
别名	TRIM62;DEAR1

## 产品验证



Western blot analysis of TRIM62 expressed in A-431, Mouse brain using TRIM62 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)