

UCKL1 Rabbit pAb

货号: **AYP18176**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	44kDa/45kDa/59kDa/61kDa
实测分子量	61kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse heart,Mouse lung,Rat pancreas
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human UCKL1 (NP_060329.2).
序列	MAAPPARADADPSPTSPTARDTPGRQAEKSETACEDRSNAESLDRLPPVGTGRSPRKRRTTSQCKSEPPLLRTSKRTIYT AGRPPWYNEHGTQSKEAFAIGLGGGSASGKTTVARMIIIEALDVPWVLLSMDSFYKVLTEQQQEAAHNNFNFDHPDAF DFDLIISTLKKLKQGKSVKVIYDFTTHSRKKDWKTYGA

靶点信息

研究背景	The protein encoded by this gene is a uridine kinase. Uridine kinases catalyze the phosphorylation of uridine to uridine monophosphate. This protein has been shown to bind to Epstein-Barr nuclear antigen 3 as well as natural killer lytic-associated molecule. Ubiquitination of this protein is enhanced by the presence of natural killer lytic-associated molecule. In addition, protein levels decrease in the presence of natural killer lytic-associated molecule, suggesting that association with natural killer lytic-associated molecule results in ubiquitination and subsequent degradation of this protein. Alternative splicing results in multiple transcript variants.
基因ID	54963
基因名	UCKL1
Swiss	Q9NWZ5
别名	UCKL1;UCK1L;URKL1

产品验证



Western blot analysis of UCKL1 expressed in Mouse heart, Mouse lung, Rat pancreas using UCKL1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn