

— ABLYBIO, Help Your Research



MMADHC Rabbit pAb

货号: **AYP18204**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	32kDa
实测分子量	33kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,NIH/3T3,HeLa,MCF7
细胞定位	Cytoplasm,Mitochondrion
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-296 of human MMADHC (NP_056517.1).
------	--------------------------------------------------------------------------------------------------------------------

靶点信息

研究背景	This gene encodes a mitochondrial protein that is involved in an early step of vitamin B12 metabolism. Vitamin B12 (cobalamin) is essential for normal development and survival in humans. Mutations in this gene cause methylmalonic aciduria and homocystinuria type cblD (MMADHC), a disorder of cobalamin metabolism that is characterized by decreased levels of the coenzymes adenosylcobalamin and methylcobalamin. Pseudogenes have been identified on chromosomes 11 and X.
基因ID	27249
基因名	MMADHC
Swiss	Q9H3L0 (https://www.uniprot.org/uniprotkb/Q9H3L0/entry)
别名	MMADHC,C2orf25,CL25022,cblD,cblD type,MMADHC Rabbit pAb,Methylmalonic aciduria and homocystinuria type D protein,mitochondrial

产品验证



Western blot analysis of MMADHC expressed in 293T,NIH/3T3,HeLa using MMADHC Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)