

# RAI14 Rabbit pAb

货号: **AYP18209**

## 产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	106kDa/109kDa/110kDa
实测分子量	130kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse lung,Rat lung
细胞定位	Cytoplasm,Nucleus,cell cortex,cytoskeleton
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 300-600 of human RAI14 (NP_056392.2).
序列	SGKESVFFAEPPFKAEISSIRENKDRLSDSTTGADSLDISSEADQQDLLSLQAKVASLTLHNKELQDKLQAKSPKEAEDLSFDYHSTQTDLGPLGKPGETSPDSSPSVLIHSLGKSTTDNDVRIQQLEILQDLQKRLESSEAERKQLQVELQSRRAELVCLNNTSEISNSDLSQKLKETQSKYEEAMKEVLSVQKQMKLGLVSPESMDNYSHFHELRTVEEINVLKQDLQNAL EESERNKEKVRELEEKLVREKGTVIKPPVEEYEEMKSSYCSVIENMNKEKAFLFEK

## 靶点信息

研究背景	Plays a role in actin regulation at the ectoplasmic specialization, a type of cell junction specific to testis. Important for establishment of sperm polarity and normal spermatid adhesion. May also promote integrity of Sertoli cell tight junctions at the blood-testis barrier.
基因ID	26064

基因名	RAI14
Swiss	Q9P0K7
别名	RAI14;NORPEG;RAI13

## 产品验证



Western blot analysis of RAI14 expressed in Mouse lung,Rat lung using RAI14 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)