

# NBEAL2 Rabbit pAb

货号: **AYP18218**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	282kDa/299kDa/302kDa
实测分子量	302kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Endoplasmic reticulum
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1800-2100 of human NBEAL2 (NP_055990.1).
序列	LLHWGALWRQLASPCGAWALRDTPIPRWKLSAETYSRMRLKLVPNHHFDPHLEASALRDNLGEVPLTPTEEASLPLAVT KEAKVSTPPELLQEDQLGEDELAELTPMEEAELDEQREKLVLSAECQLVTVVAVVPGLLLEVTTQNVYFYDGGSTERVETEEG IGYDFRRPLAQLREVHLRRFNLRRSALELFFIDQANYFLNFPCKVGTTPVSSPSQTPRPQPGPIPPHTQVRNQVYSWLLRLR PPSQGYLSSRSPQEMLRASGLTQKWVQREISNFEYLMQLNTIAGRITYNDLSQYPVFP

## 靶点信息

研究背景	The protein encoded by this gene contains a beige and Chediak-Higashi (BEACH) domain and multiple W D40 domains, and may play a role in megakaryocyte alpha-granule biogenesis. Mutations in this gene are a cause of gray platelet syndrome.
基因ID	23218

基因名	NBEAL2
Swiss	Q6ZNJ1
别名	NBEAL2;BDPLT4;GPS

## 产品验证



Western blot analysis of NBEAL2 expressed in HeLa using NBEAL2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)