

# USP6NL Rabbit pAb

货号: **AYP18237**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	94kDa/95kDa
实测分子量	115kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HeLa, LO2
细胞定位	Cytoplasmic vesicle, Golgi apparatus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 300-600 of human USP6NL (NP_055503.1).
序列	SYTILKHLHKHLMKLSMEELVEFFQETLAKDFFEDDFVIEQLQISMTTELKRAKLDLPEPGKEDEYPKKPLGQLPPELQSWG VHHLNSNGQRSVGRPSPLASGRRESGAPHRHEHSPHPQSRGTPERAQPPRRKSVEEESKLLKDEADFQRKLPSPGQDS SRQYNHAAANQNSNATSNIRKEFVPKWNKPSDVSATERTAKYTMEGKGRAAHPALAVTVPGPAEVRVSNVRPKMKALDA EDGKRGSTASQYDNVPGPELDSGASVEEALERAYSQSPRHLYPPSPRKHAEPSSSPSKVS

## 靶点信息

研究背景	Acts as a GTPase-activating protein for RAB5A and RAB43. Involved in receptor trafficking. In complex with EPS8 inhibits internalization of EGFR. Involved in retrograde transport from the endocytic pathway to the Golgi apparatus. Involved in the transport of Shiga toxin from early and recycling endosomes to the trans-Golgi network. Required for structural integrity of the Golgi complex.
------	---

基因ID	9712
基因名	USP6NL
Swiss	Q92738
别名	USP6NL;RNTRE;TRE2NL;USP6NL-IT1

## 产品验证



Western blot analysis of USP6NL expressed in HeLa, LO2 using USP6NL Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)