

CHST2 Rabbit pAb

货号: **AYP18248**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	52kDa/57kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG
细胞定位	Golgi apparatus,Single-pass type II membrane protein,trans-Golgi network membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 231-530 of human CHST 2 (NP_004258.2).
序列	FQLYSPAGSGGRNLTTLGIFGAATNKVVCSSPLCPAYRKEVVGLVDDRVCCKCPPQRLARFEEECRKYRTLVIKGVRFVDV AVLAPLLRDPALDLKVIHLVRDPRAVASSRIRSRHGLIRESLQVVRSDPRAHRMPFLEAAGHKLGAKEGVGGPADYHAL GAMEVICNSMAKTLQATALQPPDWLQGHYLVVRYEDLVGDPVKTLRRVYDFVGLLVSPMEQFALNMTSGSGSSSKPFVV SARNATQAANAWRTALTFFQIKQVEEFCYQPMAVLGYERVNSPEEVKDLSTLLRKPRL

靶点信息

研究背景	This locus encodes a sulfotransferase protein. The encoded enzyme catalyzes the sulfation of a nonreducing N-acetylglucosamine residue, and may play a role in biosynthesis of 6-sulfosialyl Lewis X antigen.
基因ID	9435

基因名	CHST2
Swiss	Q9Y4C5
别名	CHST2;C6ST;GST-2;GST2;Gn6ST-1;HEL-S-75;glcNAc6ST-1

产品验证



Western blot analysis of CHST2 expressed in U-251MG using CHST2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn