

— ABLYBIO, Help Your Research



KDM5C Rabbit pAb

货号: **AYP18256**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:1000 - 1:5000
理论分子量	155kDa/170kDa/175kDa
实测分子量	180KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,Mouse brain
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1400 to the C-terminus of human KD M5C (XP_011529128.1).
------	---

靶点信息

研究背景	This gene is a member of the SMCY homolog family and encodes a protein with one ARID domain, one JmjC domain, one JmjN domain and two PHD-type zinc fingers. The DNA-binding motifs suggest this protein is involved in the regulation of transcription and chromatin remodeling. Mutations in this gene have been associated with X-linked mental retardation. Alternative splicing results in multiple transcript variants.
基因ID	8242
基因名	KDM5C
Swiss	P41229 (https://www.uniprot.org/uniprotkb/P41229/entry)
别名	KDM5C,DXS1272E,JARID1C,MRX13,MRXJ,MRXSCJ,MRXSJ,SMCX,XE169,KDM5C Rabbit pAb,Histone demethylase JARID1C,Jumonji/ARID domain-containing protein 1C,Protein SmcX,Protein Xe169,[histone H3]-trimethyl-L-lysine(4) demethylase 5C

产品验证



Western blot analysis of KDM5C expressed in K-562, Mouse brain using KDM5C Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)