

# KDM5C Rabbit pAb

货号: **AYP18256**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:1000 - 1:5000
理论分子量	155kDa/170kDa/175kDa
实测分子量	180KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,Mouse brain
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1400 to the C-terminus of human KDM5C (XP_011529128.1).
序列	ERHGSRRARGRALERRRRRKVDRGGEGDDPAREELEPKRVRSSGPEAEVQEEEELEETGGEGPPAPIPTTGSPSTQENQ NGLEPAEGTTSGPSAPFSTLTPRLHLPCPQQPPQQQL

## 靶点信息

研究背景	This gene is a member of the SMCY homolog family and encodes a protein with one ARID domain, one JmjC domain, one JmjN domain and two PHD-type zinc fingers. The DNA-binding motifs suggest this protein is involved in the regulation of transcription and chromatin remodeling. Mutations in this gene have been associated with X-linked mental retardation. Alternative splicing results in multiple transcript variants.
基因ID	8242

基因名	KDM5C
Swiss	P41229
别名	KDM5C;DXS1272E;JARID1C;MRX13;MRXJ;MRXSCJ;MRXSJ;SMCX;XE169

## 产品验证



Western blot analysis of KDM5C expressed in K-562, Mouse brain using KDM5C Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)