

NSMAF Rabbit pAb

货号: **AYP18258**

产品信息

| | |
|-------|--|
| 反应 | Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 104kDa/107kDa |
| 实测分子量 | 110kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Mouse brain |
| 细胞定位 | cytoplasm, cytosol |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human NSMAF (NP_003571.2). |
| 序列 | MAFIRKKQQEQQLQLYSKERFSLLLLNLLEEYFEQHRANHILHKGSHHERKIRGSLKICSKSVIFEPDSISQPIIKIPLRDCIKIGKHGENGANRHFTKAKSGGISLIFSQVYFIKEHNVVAPYKIERGKMEYVFELDVPGKVEDVVETLLQLHRASCLDKLGDQTA MITAILQSRLARTSFDKNRFQNISEKLMHECKAEMVTPLVNPGHVCITDTNLYFQPLNGYPKPVVQITLQDVRRIYKRRHGLMPLGLEVFCTEDDLCSDIYLFYEPQDRDDLYFYIATYLEHHVAEHTAES |

靶点信息

| | |
|------|---|
| 研究背景 | This gene encodes a WD-repeat protein that binds the cytoplasmic sphingomyelinase activation domain of the 55kD tumor necrosis factor receptor. This protein is required for TNF-mediated activation of neutral sphingomyelinase and may play a role in regulating TNF-induced cellular responses such as inflammation. Alternative splicing results in multiple transcript variants. |
|------|---|

| | |
|-------|------------------|
| 基因ID | 8439 |
| 基因名 | NSMAF |
| Swiss | Q92636 |
| 别名 | NSMAF;FAN;GRAMD5 |

产品验证



Western blot analysis of NSMAF expressed in Mouse brain using NSMAF Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn