

TAF1 Rabbit pAb

货号: **AYP18265**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	174-182kDa, 204-216kDa
实测分子量	300kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,Jurakt,HeLa,NIH/3T3
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 990-1250 of human TAF 1 (NP_620278.1).
序列	GTDADLRRLSLKNAKQLLRKFGVPEEEIKLSRWEVIDVVRTMSTEQARSQEGPMSKFAFGSRFSVAEHQERYKEECQRIFDLQNKVLSSTEVLSTDDTSSSAEDSDFEEMGKNIENMLQNKKTSSQLSREREEQERKELQRMLLAAGSAASGNNHRDDDTASVTSLSNNSATGRCLKIYRTRFRDEEGKEYVRCETVRKPAVIDAYVRIKTTKDEEFIRKFALEFDEQHREEMRKERRRIQEQLRRLKRNQEKEKLGPEKK

靶点信息

研究背景	Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is the basal transcription factor TFIID, which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes the largest subunit of TFIID. This subunit binds to core promoter sequences encompassing the transcription start site. It also binds to activators and other transcriptional regulators, and these interactions affect the rate of transcription initiation. This subunit contains two independent protein kinase domains at the N- and C-terminals, but also possesses acetyltransferase activity and can act as a ubiquitin-activating/conjugating enzyme. Alternative splicing of this gene results in multiple transcript variants. This gene is part of a complex transcription unit (TAF1/DYT3), wherein some transcript variants share exons with TAF1 as well as additional downstream DYT3 exons.
基因ID	6872
基因名	TAF1
Swiss	P21675
别名	TAF1;BA2R;CCG1;CCGS;DYT3;DYT3/TAF1;KAT4;MRXS33;N-TAF1;NSCL2;OF;P250;TAF(II)250;TAF2A;TAFII-250;TAFII250;XDP

产品验证



Western blot analysis of TAF1 expressed in U-87MG, Jurkat, HeLa using TAF1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn