

# MAN2C1 Rabbit pAb

货号: **AYP18288**

## 产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	104kDa/113kDa/115kDa/117kDa
实测分子量	128kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney,Mouse liver
细胞定位	cytosol,nucleoplasm
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human MAN2C1 (NP_006706.2).
序列	MAAAPALKHWRTTLERVEKVFVSPPLYFTDCNLRGRLFGASCPVAVLSSFLTPERLPYQEAVQRDFRPAQVGDSFGPTWWT CWFRVELTIPEAWVGQEVHLCWESDGEGLVWRDGEVQGLTKEGKTSYVLTDRLGERDPRSLTYEVACNGLLGAGK GSMIAAPDPEKMFQLSRAELAVFHRDVHMLLVLDLELLGLIAKGLGKDNQRSFQALYTANQMVNVCDPAQPETFPVAQALA SRFFGQHGGESQHTIHATGHCHIDTAWLWPFKETVRKCARSWVTALQLMERNPEFIFACSQA

## 靶点信息

研究背景	Cleaves alpha 1,2-, alpha 1,3-, and alpha 1,6-linked mannose residues from glycoproteins. Involved in the degradation of free oligosaccharides in the cytoplasm.
基因ID	4123

基因名	MAN2C1
Swiss	Q9NTJ4
别名	MAN2C1;MAN6A8;MANA;MANA1

## 产品验证



Western blot analysis of MAN2C1 expressed in Mouse kidney, Mouse liver using MAN2C1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)