

# EXTL2 Rabbit pAb

货号: **AYP18299**

## 产品信息

|       |  |
|-------|--|
| 反应    | Human,Mouse  |
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  |  |
| 应用    | WB   |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000  |
| 理论分子量 | 37kDa  |
| 实测分子量 | 37kDa  |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  | HeLa,Mouse heart   |
| 细胞定位  | Endoplasmic reticulum membrane,Secreted,Single-pass type II membrane protein                         |
| 纯化    | Affinity purification  |

## 抗原信息


|      |  |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 41-330 of human EXTL2 (NP_001430.1).   |
| 序列   | PSVKEDKMLMLRREIKSQGKSTMDSFTLIMQTYNRTDLLLKLLNHYQAVPNLHKVIVVWNNIGEKAPDELWNSLGPHPIPIVIFKQQTANRMRNRLQVFPELETNAVLVMDLSTLSTPDLVFAFSVWQQFPDQIVGFVPRKHVSTSSGIYSYGSFEMQAPGSGNGDQYSMVLIGASFFNSKYLELFQRQPAAVHALIDDTQNCDDIAMNFIIAKHIGKTSQIFVKPVNMDNLEKETNSGYSGMWHRAEHALQRSYCINKLVNIYDSMPLRYSNIMISQGFYANYKRKI |

## 靶点信息

|      |  |
|------|--|
| 研究背景 | Glycosyltransferase required for the biosynthesis of heparan-sulfate and responsible for the alternating addition of beta-1-4-linked glucuronic acid (GlcA and alpha-1-4-linked N-acetylglucosamine (GlcNAc units to nascent heparan sulfate chains. |
| 基因ID | 2135   |

|       |             |
|-------|-------------|
| 基因名   | EXTL2       |
| Swiss | Q9UBQ6      |
| 别名    | EXTL2;EXTR2 |

## 产品验证



Western blot analysis of EXTL2 expressed in HeLa, Mouse heart using EXTL2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)