

CSNK1D Rabbit pAb

货号: **AYP18304**

产品信息

反应	Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	46kDa/47kDa
实测分子量	47kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Rat heart, Rat brain
细胞定位	Cell membrane, Cytoplasm, Golgi apparatus, Nucleus, centrosome, cytoskeleton, microtubule organizing center, perinuclear region, spindle
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 250-350 of human CSNK1D (NP_620693.1).
序列	TYLNFCRSLRFDDKPDYSYLRQLFRNLFRHQGFSYDYVFDWNMLKFGASRAADDAERERRDREERLRHSRNPATRGLPS TASGRLRGTQEVAPPTPLTPTS

靶点信息

研究背景	This gene is a member of the casein kinase I (CKI) gene family whose members have been implicated in the control of cytoplasmic and nuclear processes, including DNA replication and repair. The encoded protein may also be involved in the regulation of apoptosis, circadian rhythm, microtubule dynamics, chromosome segregation, and p53-mediated effects on growth. The encoded protein is highly similar to the mouse and rat CK1 delta homologs. Three transcript variants encoding different isoforms have been found for this gene.
------	---

基因ID	1453
基因名	CSNK1D
Swiss	P48730
别名	CSNK1D;ASPS;CKIdelta;FASPS2;HCKID

产品验证



Western blot analysis of CSNK1D expressed in Rat heart,Rat brain using CSNK1D Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn