

— ABLYBIO, Help Your Research



TRAF7 Rabbit pAb

货号: **AYP18321**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	66kDa/74kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HT-29, HeLa
细胞定位	Cytoplasmic vesicle
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 260-400 of human TRAF 7 (NP_115647.2).
------	--

靶点信息

研究背景	Tumor necrosis factor (TNF; see MIM 191160) receptor-associated factors, such as TRAF7, are signal transducers for members of the TNF receptor superfamily (see MIM 191190). TRAFs are composed of an N-terminal cysteine/histidine-rich region containing zinc RING and/or zinc finger motifs; a coiled-coil (leucine zipper) motif; and a homologous region that defines the TRAF family, the TRAF domain, which is involved in self-association and receptor binding.
基因ID	84231
基因名	TRAF7
Swiss	Q6Q0C0 (https://www.uniprot.org/uniprotkb/Q6Q0C0/entry)
别名	TRAF7,RFWD1,RNF119,TRAF7 Rabbit pAb,RING finger and WD repeat-containing protein 1,RING finger protein 119,RING-type E3 ubiquitin transferase TRAF7,TNF receptor-associated factor 7

产品验证



Western blot analysis of TRAF7 expressed in HT-29, HeLa using TRAF7 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)