

# TRAF7 Rabbit pAb

货号: **AYP18321**

## 产品信息

|       |  |
|-------|--|
| 反应    | Human  |
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  |  |
| 应用    | WB   |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000  |
| 理论分子量 | 66kDa/74kDa  |
| 实测分子量 | 75kDa  |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  | HT-29, HeLa  |
| 细胞定位  | Cytoplasmic vesicle  |
| 纯化    | Affinity purification  |

## 抗原信息


|      |  |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 260-400 of human TRAF 7 (NP_115647.2).                               |
| 序列   | PHSKYGCTFIGNQDTYETHLETCTRFEGFLKEFLQQTDDRFHEMHVALAQKDQEIAFLRSMLGKLSKIDQLEKSLELKFVDL<br>DENQSKLSEDLMEFRRDASMLNDELSHINARLNMGILGSYDPQQIFKCKGTFVGHQGPVW |

## 靶点信息

|      |  |
|------|--|
| 研究背景 | Tumor necrosis factor (TNF; see MIM 191160) receptor-associated factors, such as TRAF7, are signal transducers for members of the TNF receptor superfamily (see MIM 191190). TRAFs are composed of an N-terminal cysteine/histidine-rich region containing zinc RING and/or zinc finger motifs; a coiled-coil (leucine zipper) motif; and a homologous region that defines the TRAF family, the TRAF domain, which is involved in self-association and receptor binding. |
| 基因ID | 84231  |

|       |                    |
|-------|--------------------|
| 基因名   | TRAF7              |
| Swiss | Q6Q0C0             |
| 别名    | TRAF7;RFWD1;RNF119 |

## 产品验证



Western blot analysis of TRAF7 expressed in HT-29, HeLa using TRAF7 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)