

# RC3H1 Rabbit pAb

货号: **AYP18333**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:200 - 1:2000
理论分子量	124kDa/125kDa
实测分子量	126KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y
细胞定位	Cytoplasm,P-body
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 834-1133 of human RC3 H1 (NP_742068.1).
序列	AAGSVEMMNVESKGMRDQRLDLQRRRAETSDDDLIPFGDRPTVSRFGAISRTSKTIYQGAGPMQAMAPQGAPTKSINIS DYSPYGTHGGWGASPYSPHQNIPSQGHFSEERERISMSEVASHGKPLPSAEREQLRLELQLNHQISQQTQLRGLEAVSN RLVLQREANTLAGQSQPPPPPPKWPGMISSEQLSLELHQVEREIGKRTRELSMENQCSLDMKSKLNTSKQAENGQPEPQ NKVPAEDLTLTFSVDPNGSALTQENISLLSNKTSSLNLSLSEDPGGGDNNDSSQRSGVTPSSAP

## 靶点信息

研究背景	This gene encodes a protein containing RING-type and C3H1-type zinc finger motifs. The encoded protein recognizes and binds to a constitutive decay element (CDE) in the 3' UTR of mRNAs, leading to mRNA deadenylation and degradation. Alternative splicing results in multiple transcript variants.
基因ID	149041

基因名	RC3H1
Swiss	Q5TC82
别名	RC3H1;RNF198;ROQUIN;roquin-1

## 产品验证



Western blot analysis of RC3H1 expressed in SH-SY5Y using RC3H1 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)