

Zfp541 Rabbit pAb

货号: **AYP18358**

产品信息

| | |
|-------|--|
| 反应 | Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:200 - 1:2000 |
| 理论分子量 | 141kDa/148kDa |
| 实测分子量 | 170kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | mouse brain,mouse testis,rat brain,rat testis |
| 细胞定位 | Nucleus |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 108-269 of mouse Zfp541 (NP_001092747.1). |
| 序列 | ALKVPKEADEGGRATGSTRKGKRQHSSPQNPLLDCSLCGKVFSSASSLSKHLYLTHSQERKHVCKVCSKAFKRQDHLTGH MLTHQKTKPFVCIQGCSSYCDYRSLRRHYEVQHGVCIKETPPEEEAYGDPTHNHVDVANQPPPSGLRSLGPPPEARSPGS VL |

靶点信息

| | |
|------|--|
| 研究背景 | Predicted to enable transcription corepressor activity. Predicted to be involved in histone deacetylation; negative regulation of transcription, DNA-templated; and regulation of transcription by RNA polymerase II. Predicted to act upstream of or within cell differentiation and spermatogenesis. Located in nucleus. Part of histone deacetylase complex. Orthologous to human ZNF541 (zinc finger protein 541). |
| 基因ID | 666528 |

| | |
|-------|---------------|
| 基因名 | |
| Swiss | |
| 别名 | ZNF541;Zfp541 |

产品验证



Western blot analysis of Zfp541 expressed in mouse brain, rat brain, mouse testis using Zfp541 Rabbit p Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120 S.

实验步骤

访问官网浏览详情: www.ablybio.cn