

TRAPPC9 Rabbit pAb

货号: **AYP18367**

产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:200 - 1:2000
理论分子量	127kDa/128kDa/139kDa
实测分子量	150kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Mouse pancreas
细胞定位	Cytoplasm, Endoplasmic reticulum, Golgi apparatus, cis-Golgi network
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 947-1246 of human TRAPPC9 (NP_113654.4).
序列	SHVKTLEAVLNFKYSGGPGHTEGYRNLISLGLHVEVEPSVFFTRVSTLPATSTRQCHLLLDVFNSTEHELTVSTRSSEALILH AGECQRMAIQVDKFNFESEFPEKQGFANPKQLEERREARGLEIHSKLGICWRIPSLKRSGEASVEGLLNQLVLEHLQL APLQWDVLVDGQPCDREAVAACQVGDPVRLEVRLTNRSPRSVGFALTVPFQDHQNGVHNYDLHDTVSVFGSSTFYL DAVQPSGQSACLGALLFLYTGDFFLHIRFHEDSTSKELPPSWFCLPSVHVCALEAQA

靶点信息

研究背景	This gene encodes a protein that likely plays a role in NF-kappa-B signaling. Mutations in this gene have been associated with autosomal-recessive mental retardation. Alternatively spliced transcript variants have been described.
基因ID	83696

基因名	TRAPPC9
Swiss	Q96Q05
别名	TRAPPC9;IBP;IKBKBBP;MRT13;NIBP;T1;TRS120

产品验证



Western blot analysis of TRAPPC9 expressed in Mouse pancreas using TRAPPC9 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn