

KCNH6 Rabbit pAb

货号: **AYP18370**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:200 - 1:2000
理论分子量	56kDa/100kDa/109kDa
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	293T, HeLa
细胞定位	Membrane, Multi-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 70-220 of human KCNH6 (NP_110406.1).
序列	TGPNTPSSAVSRLAQALLGAECKVDILYRDKDASSFRCLVDVVPVKNEGDGAVIMFILNFEDLAQLLAKCSSRSLSQLLSQSFLGSEGSHGRPGGPGGTGRGKYRTISQIPQFTLNFVEFNLEKHRSSSTTEIEIAPHKVVERTQNV

靶点信息

研究背景	Voltage-gated potassium (Kv) channels represent the most complex class of voltage-gated ion channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume. This gene encodes a member of the potassium channel, voltage-gated, subfamily H. This member is a pore-forming (alpha) subunit. Alternative splicing results in multiple transcript variants that encode different isoforms.
------	---

基因ID	81033
基因名	KCNH6
Swiss	Q9H252
别名	KCNH6;ERG-2;ERG2;HERG2;Kv11.2;hERG-2

产品验证



Western blot analysis of KCNH6 expressed in 293T, HeLa using KCNH6 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn