

# NOC4L Rabbit pAb

货号: **AYP18381**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:200 - 1:2000
理论分子量	58kDa
实测分子量	65kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG, OVCAR-3, Jurkat
细胞定位	Multi-pass membrane protein, Nucleus, Nucleus membrane, nucleolus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-260 of human NOC4L (NP_076983.1).
序列	MEREPGAAGVRRALGRRLEAVLASRSEANAVFDILAVLQSEDQEEIQEAVRTCSRLFGALLERGFVQGLPSEEMVMTGS QGATRKYKVVWMRHRYSCCNRLGELLGHPSFQVKELALSALLKFVQLEGAHPLEKSKWEGNYLFPRELFKLVGGLLSPE EDQSLLLSQFREYLDYDDTRYHTMQAAVDAVARVTGQHPEVPPAFWNNAFLLSAVSLPRREPTVSSFYVKRAELWDTW KVAHLKEHRRVFQAMWLSFL

## 靶点信息

研究背景	Enables RNA binding activity. Predicted to be involved in rRNA processing. Located in nucleolus and nucleoplasm.
基因ID	79050

基因名	NOC4L
Swiss	Q9BVI4
别名	NOC4L;NET49;NOC4;UTP19

## 产品验证



Western blot analysis of NOC4L expressed in U-251MG, OVCAR-3, Jurkat using NOC4L Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)