

— ABLYBIO, Help Your Research



# CDCA7L Rabbit pAb

货号: **AYP18408**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:200 - 1:2000
理论分子量	46kDa/48kDa/52kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,Jurkat,Mouse spleen,Mouse brain,Rat brain
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human CDCA7 L (NP_061189.2).
------	---

## 靶点信息

研究背景	Plays a role in transcriptional regulation as a repressor that inhibits monoamine oxidase A (MAOA activity and gene expression by binding to the promoter. Plays an important oncogenic role in mediating the full t transforming effect of MYC in medulloblastoma cells. Involved in apoptotic signaling pathways; May act do wnstream of P38-kinase and BCL-2, but upstream of CASP3/caspase-3 as well as CCND1/cyclin D1 and E2 F1.
基因ID	55536
基因名	CDCA7L
Swiss	Q96GN5 ( <a href="https://www.uniprot.org/uniprotkb/Q96GN5/entry">https://www.uniprot.org/uniprotkb/Q96GN5/entry</a> )
别名	CDCA7L,JPO2,R1,RAM2,CDCA7L Rabbit pAb,Protein JPO2,Transcription factor RAM2,HR1

## 产品验证



Western blot analysis of CDCA7L expressed in Mouse spleen,Rat brain,A-549 using CDCA7L Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)