

CDCA7L Rabbit pAb

货号: **AYP18408**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:200 - 1:2000
理论分子量	46kDa/48kDa/52kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,Jurkat,Mouse spleen,Mouse brain,Rat brain
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human CDCA7 L (NP_061189.2).
序列	MELATRYQIPKEVADIFNAPSDDEEFVGFRRDDVPMETLSSEESCDSFDSLESGKQQDVRFHSHKYFTEELRRIFIEDTSETEDFAGFTQSDLNGKTNPEVMVVEDLSDDGKASLVSEEEEEDEEDKATPRRSRSRRSSIGLRVAFQFPTKLANPKDNSSSEQLFSSARLQNEKKTILERKKDCRQVIQREDSTSESEDDSRDESQESSDALLKRTMNIKENKAMLAQLLAELNSMPDFFPVRTPTSASRKKTVRRAFSEGQITRRMNPTRSARPPEKFALENFTVSAAKFAEEFYF

靶点信息

研究背景	Plays a role in transcriptional regulation as a repressor that inhibits monoamine oxidase A (MAOA activity and gene expression by binding to the promoter. Plays an important oncogenic role in mediating the full transforming effect of MYC in medulloblastoma cells. Involved in apoptotic signaling pathways; May act downstream of P38-kinase and BCL-2, but upstream of CASP3/caspase-3 as well as CCND1/cyclin D1 and E2F1.
------	--

基因ID	55536
基因名	CDCA7L
Swiss	Q96GN5
别名	CDCA7L;JPO2;R1;RAM2

产品验证



Western blot analysis of CDCA7L expressed in Mouse spleen,Rat brain,A-549 using CDCA7L Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn