

— ABLYBIO, Help Your Research



# PXK Rabbit pAb

货号: **AYP18421**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:200 - 1:2000
理论分子量	38kDa/49kDa/55kDa/58kDa/60kDa/62kDa/64kDa
实测分子量	80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,U-251MG,Jurkat,Mouse spleen,Mouse lung,Rat kidney
细胞定位	Cell membrane,Cytoplasm,Peripheral membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human PXK (N P_060241.2).
------	--

## 靶点信息

研究背景	This gene encodes a phox (PX) domain-containing protein which may be involved in synaptic transmission and the ligand-induced internalization and degradation of epidermal growth factors. Variations in this gene may be associated with susceptibility to systemic lupus erythematosus (SLE). Alternative splicing results in multiple transcript variants.
基因ID	54899
基因名	PXK
Swiss	Q7Z7A4 ( <a href="https://www.uniprot.org/uniprotkb/Q7Z7A4/entry">https://www.uniprot.org/uniprotkb/Q7Z7A4/entry</a> )
别名	PXK,MONAKA,SLOB,PXK Rabbit pAb,Modulator of Na,K-ATPase

## 产品验证



Western blot analysis of PXK expressed in Mouse spleen, Rat kidney, HepG2 using PXK Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))