

# TACO1 Rabbit pAb

货号: **AYP18430**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IHC:</b> 1:50 - 1:200
理论分子量	32kDa
实测分子量	32kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,LO2,A375,Mouse brain,Mouse heart,Mouse kidney,Mouse liver,Rat lung
细胞定位	Mitochondrion
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-297 of human TACO1 (NP_057444.2).
序列	MSAWAAASLSRAAARCLLARGPGVRAAPPRDPRPSHPEPRGCGAAPGRTLHFTAAPVAGHNKWSKVRHIKGPKDVERSR IFSKLCLNIRLAVKEGGPNPEHNSNLANILEVCRSKHMPKSTIETALKMEKSKDITYLLYEGRGGSSLLIEALSNSHKCQA DIRHILNKNGGVMAVGARHSFDKKGIVVEVEDREKKAVNLERALEMAIEAGAEDVKETEDEEERNVFKFICDASSLHQVR KKLDSLGLCSVSCALEFIPNSKVQLAEPDLEQAAHLIQALSNHEDVIHVYDNIE

## 靶点信息

研究背景	This gene encodes a mitochondrial protein that function as a translational activator of mitochondrially-en coded cytochrome c oxidase 1. Mutations in this gene are associated with Leigh syndrome.
基因ID	51204

基因名	TACO1
Swiss	Q9BSH4
别名	TACO1;CCDC44

## 产品验证



Western blot analysis of TACO1 expressed in Mouse brain,Rat lung,U-251MG using TACO1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)