

— ABLYBIO, Help Your Research



# TBL2 Rabbit pAb

货号: **AYP18442**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:200 - 1:2000
理论分子量	49kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,293T,HeLa,Mouse brain,Rat pancreas
细胞定位	endoplasmic reticulum
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 100-430 of human TBL2 (NP_036585.1).
------	--

## 靶点信息

研究背景	This gene encodes a member of the beta-transducin protein family. Most proteins of the beta-transducin family are involved in regulatory functions. This protein is possibly involved in some intracellular signaling pathway. This gene is deleted in Williams-Beuren syndrome, a developmental disorder caused by deletion of multiple genes at 7q11.23.
基因ID	26608
基因名	TBL2
Swiss	Q9Y4P3 ( <a href="https://www.uniprot.org/uniprotkb/Q9Y4P3/entry">https://www.uniprot.org/uniprotkb/Q9Y4P3/entry</a> )
别名	TBL2,WBSCR13,WS-betaTRP,TBL2 Rabbit pAb,WS beta-transducin repeats protein,Williams-Beuren syndrome chromosomal region 13 protein

## 产品验证



Western blot analysis of TBL2 expressed in Mouse brain,Rat pancreas,LO2 using TBL2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)