

EDC4 Rabbit pAb

货号: **AYP18451**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:200 - 1:2000
理论分子量	109kDa/151kDa
实测分子量	160-180kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,Mouse liver
细胞定位	Cytoplasm,Nucleus,P-body
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 800-1100 of human EDC 4 (NP_055144.3).
序列	GDGDRHNTPSLLEAALTQEASTPDSQVWPTAPDITRETCSTLAESPRNGLQEKHKSLAFHRPPYHLLQQRDSQDASAEQSDHDDEVASLASASGGFGTKVPAPRLPAKDWKTKGSPRTSPKLRKSKKDDGDAAMGSRLTEHQVAEPPEDWPALIWQQRELAELRHSQEELLQRLCTQLEGLQSTVTGHVERALETRHEQEQRRLERALAEGQQRGGQLQEQLTQQLSQALSSAVAGRLERSIRDEIKKTVPCVSRSLPEMAGQLSNSVATKLTAVEGSMKENISKLLKSKNLTDIAIR

靶点信息

研究背景	In the process of mRNA degradation, seems to play a role in mRNA decapping. Component of a complex containing DCP2 and DCP1A which functions in decapping of ARE-containing mRNAs. Promotes complex formation between DCP1A and DCP2. Enhances the catalytic activity of DCP2 (in vitro).
基因ID	23644

基因名	EDC4
Swiss	Q6P2E9
别名	EDC4;GE1;Ge-1;HEDL5;HEDLS;RCD-8;RCD8

产品验证



Western blot analysis of EDC4 expressed in Jurkat, Mouse liver using EDC4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5 % non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn