

STK38L Rabbit pAb

货号: **AYP18454**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:200 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	43kDa/54kDa
实测分子量	54kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Mouse small intestine,Mouse brain,Rat stomach,Rat brain
细胞定位	Cytoplasm,Membrane,cytoskeleton
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 165-464 of human STK38L (NP_055815.1).
序列	LIMEFLPGGDMMTLLMKKDTLTEEETQFYISETVLAIDAIHQLGFIHRDIKPDNLLDDAKGHVKLSDFGLCTGLKKAHRTEFY RNLTHNPPSDFSFQNMNSKRKAETWKKNRRQLAYSTVGTPDYIAPEVFMQTGYNKLCDDWWSLGVIMYEMLIGYPPFCSE TPQETYRKVMNWKETLVFPPEVPISEKAKDLILRFCIDSENRIKNSGVVEIKGHPFFEGVDWEHIRERPAAPIEIKSIDDTSNF DDFPESDILQVPNTTEPDYKSKDWWFLNYTYKRFEGLTQRGSIPTYMKAGKL

靶点信息

研究背景	Involved in the regulation of structural processes in differentiating and mature neuronal cells.
基因ID	23012

基因名	STK38L
Swiss	Q9Y2H1
别名	STK38L;NDR2

产品验证



Western blot analysis of STK38L expressed in Mouse small intestine, Rat stomach, HeLa using STK38L Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30 ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 1 20s.

实验步骤

访问官网浏览详情: www.ablybio.cn