

# WDHD1 Rabbit pAb

货号: **AYP18458**

## 产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:200 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	112kDa/125kDa
实测分子量	135kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,U-251MG,A-431,HeLa,Jurkat
细胞定位	Nucleus,nucleoplasm
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 830-1129 of human WD HD1 (NP_009017.1).
序列	EEEEEDFRKKNAGYSNTATEWSQPRFRNQVEEDAEDSGEADDEEKPEIHKPGQNSFSKSTNSSDVSASGAVTFSSQ GRVNPFKVSASSKEPAMSMNSARSTNILDNMGKSSKSTALSRTTNEKSPIIKPLIPKPKPKQASAASYFQKRNSQTNKT EEVKEENLKNVLSETPAICPPQNTENQRPKTGFQMWLEENRSNILSDNPDFSDEADIIKEGMIRFRVLSTEERKVVANKAK GETASEGTEAKKRKRVDSEDETENQEEKAKENLNLSKKQKPLDFSTNQKLSAFKQE

## 靶点信息

研究背景	The protein encoded by this gene contains multiple N-terminal WD40 domains and a C-terminal high mobility group (HMG) box. WD40 domains are found in a variety of eukaryotic proteins and may function as a daptor/regulatory modules in signal transduction, pre-mRNA processing and cytoskeleton assembly. HMG boxes are found in many eukaryotic proteins involved in chromatin assembly, transcription and replication. Alternative splicing results in two transcript variants encoding different isoforms.
基因ID	11169
基因名	WDHD1
Swiss	O75717
别名	WDHD1;AND-1;AND1;CHTF4;CTF4

## 产品验证



Western blot analysis of WDHD1 expressed in LO2,U-251MG,A-431 using WDHD1 Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)