

# FZD10 Rabbit pAb

货号: **AYP18460**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:200 - 1:2000
理论分子量	65kDa
实测分子量	72kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	A-549, HeLa
细胞定位	Cell membrane, Multi-pass membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 135-230 of human FZD10 (NP_009128.1).
序列	RKLPNKNDPNYLCMEAPNNGSDEPTRGSLFPPLFRPQRPHSAQEHLKDGGPGRGGCDNPGKFHHVEKSASCAPLCTPGVDVYWSREDKRFVAVVW

## 靶点信息

研究背景	This gene is a member of the frizzled gene family. Members of this family encode 7-transmembrane domain proteins that are receptors for the Wingless type MMTV integration site family of signaling proteins. Most frizzled receptors are coupled to the beta-catenin canonical signaling pathway. Using array analysis, expression of this intronless gene is significantly up-regulated in two cases of primary colon cancer.
基因ID	11211

基因名	FZD10
Swiss	Q9ULW2
别名	FZD10;CD350;FZ-10;Fz10;FzE7;hFz10

## 产品验证



Western blot analysis of FZD10 expressed in A-549, HeLa using FZD10 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)